ABSTRAK

Waira Murib, Universitas Indonesia Timur, Epidemiologi Faktor Risiko Demam Typoid Di Puskesmas Bhintuka (Dibimbing oleh Rosdiana R dan Marthini)

Demam Tifoid Adalah Penyakit Infeksi Yang Disebabkan Oleh Salmonella Typhi Yang Biasanya Terdapat Pada Saluran Pencernaan Dengan Gejala Demam Berkepanjangan, Gangguan Saluran Pencernaan Dan Gangguan Kesadaran. Data Who (World Health Organisation) Memperkirakan Angka Insidensi Di Seluruh Dunia Terdapat Sekitar 17 Juta Pertahun Dengan 600.000 Orang Meninggal Karena Demam Typoid Dan 70% Kematianya Terjadi Di Asia

Penelitian Ini Bertujuan Untuk Mengetahui Faktor Risiko Demam Typoid. Desain Pelitian Yang Digunakan Adalah *Case Control Study* Dengan Teknik Pengambilan Sampel Secara *Simple Random Sampling* Yang Terdiri Dari 68 Kasus Dan 68 Kontrol.

Hasil Analisis *Odss Ratio* Dengan *Confidenc Interval* (Ci) = 95%Menunjukkan Bahwa Umur Merupakan Faktor Risiko Terhadap Demam Typoid Dengan Nilai Or = 35,00, Pendidikan Merupakan Faktor Risiko Terhadap Demam Typoid Dengan Nilai Or = 36,52, Spal Merupakan Faktor Risiko Terhadap Demam Typoid Dengan Nilai Or = 8,38 Dan Personal Hygiene Merupakan Faktor Risiko Terhadap Demam Typoid Dengan Nilai Or = 23,16.

Diharapkan Perlunya Perbaikan Kuantitas Dan Kualitas Sanitasi Lingkungan Seperti Air Limbah, Perlunya Penyuluhan Yang Berkesinambungan Mengenai Higene Perorangan. Perlunya Penelitian Lanjutan Mengenai Faktor Risiko Lain Yang Berhubungan Dengan Kejadian Demam Tifoid.

Kata Kunci : Umur, Tingkat Pendidikan, Spal, Personal Hygiene Dan Demam Typoid.

Daftar Pustaka : (1990 – 2014)

ABSTRACT

Waira Murib, University Of East Indonesia, Epidemiology Risk Factors For Typhoid Fever At The Bhintuka Health Center *(Supervised By Mrs. Rosdiana R dan Mrs. Marthini)*

Typhoid Fever Is An Infectious Disease Caused By Salmonella Typhi Which Is Usually Found In The Digestive Tract With Symptoms Of Prolonged Fever, Digestive Tract Disorders And Impaired Consciousness. Who (World Health Organization) Data Estimates That The Incidence Rate Worldwide Is Around 17 Million Per Year With 600,000 People Dying From Typhoid Fever And 70% Of Deaths Occurring In Asia.

This Study Aims To Determine The Risk Factors For Typhoid Fever. The Research Design Used Was A Case-Control Study With A Simple Random Sampling Technique Consisting Of 68 Cases And 68 Controls.

The Results Of The Odss Ratio Analysis With A Confidence Interval (Ci) = 95% Indicate That Age Is A Risk Factor For Typhoid Fever With An Or Value Of 35.00, Education Is A Risk Factor For Typhoid Fever With An Or Value Of 36.52, Spal Is A Risk Factor For Typhoid Fever. Typhoid Fever With An Or Value Of 8.38 And Personal Hygiene Is A Risk Factor For Typhoid Fever With An Or Value Of 23.16.

The Need To Increase The Quantity And Quality Of Environmental Sanitation Such As Waste Water, The Need For Counseling On Matters Concerning Individuals. The Need For Further Research On Other Risk Factors Associated With The Incidence Of Typhoid Fever.

Keywords *: Age, Education Level, Spal, Personal Hygiene And Typhoid Fever*

Bibliography *: (1990 – 2014)*